



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

RHYNCHITIDÆ.

Rhynchites bicolor *Fab.* S., S.

OTIORHYNCHIDÆ.

Thricolepis inornata *Horn.* F., C.

CURCULIONIDÆ.

Sitones sp.

Apion sp.

Stephanocleonus plumbeus *Lec.* F., C.

Otidocephalus estratus *Csy.* F., S.

Anthonomus decipiens *Lec.* F., S.

Tyloderma baridium *Lec.* F., C.

Ceutorhynchus pusio *Mann.* F., S.

CALANDRIDÆ.

Cossonus subareatus *Boh.* F., C.

concinus *Boh.* S., S.

SCOLYTIDÆ.

Gnatatrichius sulcatus. H., S.

Tomicus calligraphus *Germ.* H., S.

plastographus *Lec.* H., S.

rectus *Lec.* H., S.

Hylastes n. sp. H., S.

n. sp. H., S.

Hylurgops subcostulatus *Mann.* H., S.

Scolytus præceps *Lec.* H., S.

Dendroctonus valens H., S.

Polygraphus rufipennis *Kby.* H., S.

ORTHOPTERA.

BY J. A. G. REHN.

PHASMIDÆ.

Diapheromera sp.

One immature female; August 17, 1901.

ACRIDIDÆ.

Stenobothrus curtipennis (Harris).

One male; August 17, 1901.

This species has previously been recorded from Utah Co., Utah, and several localities in Colorado (altitudes 5,000–8,500 feet), the most southern being Garland, Costillo County.

Arphia arcta Scudder.

One female; August 17, 1901.

Camnula pellucida (Scudder).

Six specimens, four males, two females; August 17, 1901.

Dissosteira carolina (Linnaeus).

One male; August 17, 1901.

Circotettix suffusus (Scudder).

One male; August 17, 1901.

Previous reports included the following Colorado records: Manitou 6,300 feet, Florrisant 8,000 feet, Alma 10,000 feet, and North Fork of South Platte River.

Melanoplus bivittatus (Say).

Ten specimens, three males, five females, two immature specimens; August 17, 1901; Denver, Colorado, August 26, 1901.

TETTIGONIDÆ.

Scudderia furcata Brunner.

One male; August 17, 1901.

Orchelimum sp.

Three specimens, one male, two females; August 17, 1901.

Ceuthophilus uniformis Scudder.

Two males; August 17, 1901.

Previous records from this general region: Plains of Northern New Mexico (eastern slope), Beaver Brook, Colorado (6,000 feet), and Manitou, Colorado (6,300 feet).

Additional Orthoptera determined by S. H. Scudder and collected by T. D. A. Cockerell, W. P. Cockerell and M. D. Cockerell.

Melanoplus cockerelli. C.

bruneri. C.

gillettei. W. P. C.

Ceuthophilus valgus. C.

Bacillus coloradus. W. P. C.

Anabrus coloradus. W. P. C.

Melanoplus bivittatus. M. D. C.

NEUROPTERA—Odonata.

BY P. P. CALVERT.

Argia sp. (new?)

1 ♀, allied to *mæsta* Hag. and *violacea* Hag.

Lestes disjunctus Selys.

4 ♂, 2 ♀. Ranges from Nova Scotia to Washington, and in the West south through Colorado.

Enallagma calverti Morse.

6 ♂. A male was taken by Prof. Cockerell at Top Range, N. Mex., 11,000 feet, on June 29. This species is known from Maine, Massachusetts and Indiana in the East. In the West it ranges from Alaska and Washington to New Mexico.